


# Calving Ease

January 2013

By Sam Leadley of Attica Veterinary Associates



## High Coliform Count in Colostrum? What to do next.

- High coliform bacteria counts in colostrum should not be normal.
- It is possible to feed colostrum with low coliform bacteria counts.
- Efforts to reduce coliform bacteria counts in colostrum must be a team effort – everyone has to buy into the goal of clean colostrum.
- It is most cost effective to focus on key critical control points:
  1. Clean teats on fresh cows
  2. Clean collection equipment
  3. Feed quickly or cool rapidly for stored colostrum
  4. Clean feeding equipment
- Monitor, monitor, monitor with sampling and lab cultures.

Yes, it is discouraging to get the lab culture results back and find high coliform counts in the colostrum. However, there is good news! It is possible to deliver clean colostrum to calves.

Let's look at the situation for one of my clients with serious health problems in young calves. At the first sampling of colostrum we had six samples that looked like this:

Sample #1= 38,000 cfu/ml; #2= 278,000; #3= 54,000; #4= 125,000; #5= 46,000; #6= 27,000. Remember, the threshold for acceptable colostrum for newborn calves for coliforms is 5,000 cfu/ml.

**What to do next?** On one hand, we could have spent considerable time and money on more sampling and culturing. One can serial sample from collection bucket to feeding the calf. That may pin down the most likely sources of inoculation with coliforms and where growth is taking place.

On the other hand, with high numbers like these and such a wide spread from highest (278,000) to lowest (27,000) I have to ask, "Is sampling and culturing really necessary?"

To correct the situation we chose to just jump in with both feet focusing on four critical bacteria control points. Teat preparation for fresh cows in the milking parlor was first. We met with all the milking staff that milked fresh cows. Each person was coached on a "dip-wipe-dip-wipe-scrub-teat -end" procedure. Each worker showed the parlor manager they could properly follow these rules.

One person was responsible for cleaning the stainless steel buckets used to collect the colostrum. Visual inspection showed significant areas of biofilm on the inside of all of the buckets. I made sure that there was a dry chlorinated detergent available as well as a stiff

bristle brush, a green scouring pad and rubber gloves. We reviewed the “rinse-wash-rinse-dry” cleaning protocol. [See [www.calffacts.com](http://www.calffacts.com) click on Washing Milk Containers: Protocol] The farm had lots of hot water so keeping the wash water above 120°F was not a problem. The insides of these buckets now shine brightly. Good.

The parlor manager agreed to change his handling of the colostrum buckets. He had been waiting until all the fresh cows were milked (sometimes as long as an hour) until taking the collection buckets from the parlor to the utility room. Then they would often sit there for another hour before being transported over to the calf barn. [Can you almost hear the coliform bacteria growing?] Once at the calf barn the colostrum was poured into 2-quart plastic containers (visualize plastic food containers). Then those went into a freezer for storage until needed. Often more than 30 containers of warm colostrum were added at one time. Too many opportunities for bacteria to grow, and grow, and grow.

A whole new approach to chilling colostrum is now in place. A chest-type freezer was added to the utility room. Two-liter plastic bottles of water are frozen in the freezer. As soon as each fresh cow is milked the collection bucket is brought to the utility room and two of these ice-filled bottles go into each bucket of colostrum. At the end of the milking shift all the colostrum is now transported to the calf barn. By the time the colostrum gets to the calf barn it now is chilled to around 60°F (16°C). [Coliform bacteria take 150 minutes to double at 60° compared to 20 minutes to double at cow body temperature, 102°.]

All the workers at the calf barn got together for training on cleaning storage and feeding equipment. Previous procedures did not include rinsing before washing. Nor were they monitoring wash water temperature – it was about 90°F the day I was there. They were using dish-washing detergent without any chlorine – not good if you want to remove milk protein.

The first change made to improve sanitation was to get the Washing Milk Containers Protocol translated into their native language. Then a native speaker explained the reason for all four steps. The success of this sanitation program was initially monitored by culturing sterile-water samples collected from the colostrum feeding equipment. [See [www.calffacts.com](http://www.calffacts.com) click on “Bacteria quality control: collecting rinse samples.”] Now, each month five “as-fed” samples of colostrum are cultured to monitor the success of the entire bacteria management program. The most recent results were all below 5,000 cfu/ml.

Remember, too, when using colostrum replacers this mixing and feeding equipment sanitation is a critical step in maintaining good calf health.

If you know of someone that doesn't currently receive **Calving Ease** but would like to, tell them to **WRITE** to Calving Ease, 11047 River Road, Pavilion, NY 14525 or to **CALL** 585-591-2660 (Attica Vet Assoc. office) or **FAX** (585-591-2898) or **e-mail** [calvingease@rochester.rr.com](mailto:calvingease@rochester.rr.com).  
A limited number of back issues may be accessed on the Internet at either [www.atticacows.com](http://www.atticacows.com) or [www.calnotes.com](http://www.calnotes.com) and clicking on the link, Calving Ease.

Check out the new Blog: Calves with Sam

Find it at:

<https://www.blogger.com/blogger.g?blogID=8405521823962376971&pli=1#allposts>

2013 DCHA Conference – April 4<sup>th</sup>-5<sup>th</sup> ~  
Lancaster, PA

<http://www.calfandheifer.org/?page=2013Conference>